

International Journal of Modern Physics A
 © World Scientific Publishing Company

A Review of Pentaquark Calculations on the Lattice

Keh-Fei Liu

Dept. of Physics and Astronomy, University of Kentucky, Lexington, KY, 40506, USA

Nilmani Mathur

Jefferson Lab., 12000 Jefferson Ave., Newport News, VA 23606, USA

We review lattice calculations of pentaquarks and discuss issues pertaining to interpolation fields, distinguishing the signal of pentaquarks from those of the KN scattering states, chiral symmetry, and ghost state contaminations.

Keywords: Pentaquarks; lattice calculation

1. Introduction

The recent interest in pentaquark baryons has been inspired by the experimental discovery of $\Theta^+(1540)$ whose quark composition is $uudd\bar{s}$. We will not address the experimental situation. It is summarized by T. Nakano in his plenary talk during this conference ¹.

In the past few years, there has been quite an effort to calculate these pentaquark states to see if they can be observed in lattice QCD calculations and if their masses and other properties agree with those observed experimentally. In particular, it would be interesting to see if there is a way to understand why the observed width of $\Theta^+(1540)$ is much narrower than those of the ordinary baryons which are of the order of several hundred MeV.

Somewhat similar to the experimental situation, among the dozen or so lattice calculations, 5 calculations ^{2,3,7,9,11} claim to have positive signals for the pentaquark; while 7 calculations ^{4,5,6,8,10,12,13} reported null results. Prior to pentaquarks, lattice calculations dwell mostly on stable particles in strong decays, such as the nucleon, the pion, and kaon. Since the quark mass in the present lattice calculations is still not light enough, the Δ , ρ , and ϕ are also below the decay thresholds and can be calculated as the ground states. Come pentaquark, the interpolation field involves two u quarks, two d quarks, and one strange antiquark which inevitably will couple to the KN scattering states in addition to the potential pentaquark state. This presents a challenge not confronted before and, presumably, posted some confusion in the first round of lattice calculations. We shall address some of the issues due to this complication, namely the questions of interpolation fields, the issue of KN scattering states, the importance of chiral symmetry, and the contamination

2 *Keh-Fei Liu*

of ghost states - a quenched artifact and will attempt to draw some lessons from these calculations.

2. Interpolation Fields

There has been quite a lot of discussion in the literature on the choice of interpolation fields. The most naive one is the product of nucleon and kaon interpolation fields, i.e.

$$\chi_1^{I=0,1} = \epsilon^{abc} (u^{Ta} C \gamma_5 d^b) [u^c (\bar{s}^e \gamma_5 d^e) \mp \{u \leftrightarrow d\}], \quad (1)$$

where the $\mp$ combination is for $I = 0$ and $I = 1$ and has been denoted as such.. Another one is similar, except with the color indices e and c positioned differently

$$\chi_2^{I=0,1} = \epsilon^{abc} (u^{Ta} C \gamma_5 d^b) [u^e (\bar{s}^e \gamma_5 d^c) \mp \{u \leftrightarrow d\}], \quad (2)$$

The third one which has been used in the lattice pentaquark calculation is

$$\chi_3^\Gamma = \epsilon^{gce} \epsilon^{ghf} \epsilon^{abc} (u^{Ta} C \gamma_5 d^b) (u^{Tf} C \Gamma d^h) \Gamma C^{-1} \bar{s}^{Te}, \quad (3)$$

with $\Gamma = \{S, A\} \equiv \{1, \gamma_\mu \gamma_5\}$ ³. It is motivated by the diquark-diquark-antiquark picture of Jaffe and Wilczek¹⁵.

The masses and spectral weights due to an interpolation field operator O can be extracted from the zero-momentum correlation function with a point source, which is a sum of exponentials for the spectrum of states with the quantum numbers of the operator O

$$\langle \sum_{\vec{x}} O(\vec{x}, t) O(\vec{x}_0, 0) \rangle = \sum_i W_i e^{-m_i t}. \quad (4)$$

It is often asserted in the literature that χ_1 , being the KN interpolation field, will couple to the KN scattering states stronger than do χ_2 and χ_3 . On the other hand, χ_3 is expected to couple weakly to KN states and more strongly to the pentaquark state. However, it has been learned in the lattice community over the years not to be overly sanguine about predicting the structure of the hadron based on the interpolation field. It is already shown in Ref. [4] that χ_1, χ_2 , and χ_3 are linearly related through the multiplication of γ_5 and a Fiertz transform between the u and $\bar{s}$ fields, i.e.

$$\gamma_5 \times (\chi_1^{I=0} - \chi_2^{I=0}) = \frac{1}{2} (\chi_3^S + \chi_3^A). \quad (5)$$

Thus they are expected to couple to the same states ($1/2^-$ and $1/2^+$) with comparable spectral weights.

As a further check, we show in Fig. 1 the effective mass plot of the ground states as obtained with the three operators in the $1/2^-$ channel for the u/d quark mass corresponding to a pion mass at 633 MeV and the strange mass corresponding to the physical ϕ mass on a $16^3 \times 28$ lattice with overlap fermions and a lattice




Fig. 1. The effective mass of the $1/2^-$ ground states from the χ_1, χ_2, χ_3 and the KN open jaw diagram as a function of time separation.




Fig. 2. The spectral weights (in arbitrary units) of the $1/2^-$ (left panel) and $1/2^+$ (right panel) ground states from the χ_1, χ_2 , and χ_3 interpolation fields.

spacing at 0.2 fm^4 . We see that at short time separation, i.e. $t \leq 7 (1.4\text{fm})$, the three interpolation fields do not give the same mass. Only when $t \sim 12 (2.4\text{fm})$ do they come down to the KN threshold. Whereas the ‘open-jaw diagram’, where the kaon and nucleon are separately projected to the zero momentum states, has better overlap with the threshold NK scattering state and come down to the threshold much earlier. This shows that the three interpolation fields project to the same ground state at large time separation. We also plot their ground state spectral weights in the $1/2^-$ and the $1/2^+$ channels in Fig. 2. We see that the spectral weight of χ_3 is somewhat smaller than those of χ_1 and χ in the $1/2^-$ channel (KN in S-wave), but not orders of magnitude smaller; whereas, all three give the same weights in the $1/2^+$ channel (KN in P-wave). A single channel approach with enough time separation will be able to determine the ground state reliably. But if the pentaquark state is close to the KN scattering state, as is considered in

several calculations as a possibility in the $1/2^-$ channel, variational approach with multiple interpolation fields is a more effective approach. As a rule of thumb, the highest state will always be contaminated by still higher states and, thus, cannot be trusted. It is with the multi-operator variational calculations^{10,13} that one realizes that the earlier signal² of a low-lying $1/2^-$ state in addition to the KN state at the threshold is spurious and is due to the fact that there were only two operators in the variational calculation.

It is also widely speculated that the diquark-diquark-antiquark interpolation field proposed by Jaffe and Wilczek¹⁵

$$\chi_{JW} = \epsilon^{gfc} \epsilon^{def} \epsilon^{abc} (u^{Ta} C \gamma_5 d^b) \overleftrightarrow{\mathcal{D}}_\mu (u^{Td} C \gamma_5 d^e) \gamma_\mu \gamma_5 C \bar{s}^{Tg} \quad (6)$$

will have a better overlap with the pentaquark state to reflect the structure of the pentaquark. We note that this operator is very similar to the operator χ_3^A

$$\chi_3^A = \epsilon^{gfc} \epsilon^{def} \epsilon^{abc} (u^{Ta} C \gamma_5 d^b) (u^{Td} C \gamma_\mu \gamma_5 d^e) \gamma_\mu \gamma_5 C \bar{s}^{Tg}. \quad (7)$$

The only difference is that there is a covariant derivative between the two diquarks in χ_{JW} , while χ_3^A has a γ_μ between u^T and d in one of the $u-d$ diquark pair.

It turns out the pentaquark correlation functions built from these two interpolation fields are linearly related. To see this, we consider the part of the correlator where the $\overleftrightarrow{\mathcal{D}}_\mu$ and γ_μ operate on the d quark propagator to the right which can be written in terms of the eigenstates of the Dirac operator, i.e. $S(x, y) = \sum_\alpha \psi_\alpha(x) \psi_\alpha^\dagger(y) / (i\lambda_\alpha + m)$. Now, the $\mathcal{D}_\mu$ and γ_μ operating on each eigenstate give

$$\mathcal{D}_j \psi_\alpha = \begin{pmatrix} \mathcal{D}_j U_\alpha \\ \mathcal{D}_j L_\alpha \end{pmatrix}; \quad \gamma_j \psi_\alpha = \begin{pmatrix} -i\sigma_j L_\alpha \\ i\sigma_j U_\alpha \end{pmatrix}. \quad (8)$$

The upper component of $\mathcal{D}_j \psi_\alpha$ and $\gamma_j \psi_\alpha$, i. e. $\mathcal{D}_j U_\alpha$ and $-i\sigma_j L_\alpha$ are related through the Dirac eigenvalue equation

$$\mathcal{D}_j U_\alpha + i\epsilon_{jki} \sigma_i \mathcal{D}_k U_\alpha = (\lambda_j - i\mathcal{D}_4) \sigma_j L_\alpha. \quad (9)$$

The lower components are similarly related. Thus, as far as the role of the interpolation fields is concerned and barring special exceptions, one would expect χ_{JW} to be equivalent to χ_3^A which is in term equivalent to χ_1 and χ_2 . There is no a priori reason why one operator will preferentially project to a particular state, be it the pentaquark or the KN scattering state. We should remark that the derivative operator χ_{JW} is motivated by the nonrelativistic picture and projects to the upper component of the Dirac eigenstates in the $1/2^+$ channel; whereas, $\gamma_5 \chi_1$, $\gamma_5 \chi_2$, and χ_3 utilize the lower components. For a large quark mass, $\sigma_j L_\alpha$ is approximately equal to $\mathcal{D}_j U_\alpha / m$ from Eq. (9). In this case, the correlation function due to a local operator is $O(m^2)$ smaller than that of the derivative operator. This is the reason that the derivative operators are usually used to calculate the orbitally excited states in heavy quarkonia. To illustrate this point, we plot in Fig. 3 the spectral




Fig. 3. The spectral weights (in arbitrary units) of S_{11} and the $N\eta'$ ghost state as a function of $m_\pi^2(\text{GeV}^2)$.

weight of the S_{11} which is obtained from the lower component of the nucleon correlation function with the $1 - \gamma_4$ projection to the negative parity state¹⁶. We see that the spectral weight does go down like m^2 in the range of pion mass from 250 MeV to 350 MeV. Thus, for heavy quarks, it is better to use the derivative operator as the interpolation field and obtain the negative-parity S_{11} state from the upper component of the S_{11} correlator. For small quark masses, the lower component of the nucleon correlator works for the S_{11} state just as well. We believe the same is true with the pentaquark calculations. One should be able to reach the same five-quark states with the derivative operator, the χ_3 operator, or equivalently the χ_1 or χ_2 operators with comparable spectral weights when the quark masses are close to their physical values.

3. Distinguishing Pentaquarks from KN Scattering States

Since the 5-quark interpolation field will, in general, project to both the one-particle pentaquark states and the two-particle KN scattering states on the lattice, one needs to devise a way to distinguish them. To this end, it was advocated to study the volume dependence of the spectral weights^{4,16}. If it is a one particle state, the spectral weight for the correlator constructed with point source and zero momentum point sink has essentially no volume dependence. On the other hand, if it is a two-particle scattering state with relatively weak interaction, it is inversely proportional to the 3 volume from the normalization factor. The detailed study of such volume dependence with overlap fermions by comparing results from the $16^3 \times 28$ and $12^3 \times 28$ lattices with pion mass m_π as low as 180 MeV revealed

that the ground states of both the $1/2^-$ and $1/2^+$ channels are scattering states⁴. One needs to be careful here in view of the fact that it is more subtle to obtain the spectral weight than the mass of a state. The former is more sensitive to the fitting procedure and the length of time separation in the correlator. As it is shown recently that only when the time separation is large enough (~ 2.7 fm) and with accurate data is the volume scaling of the spectral weight for the $I = 2$ $\pi\pi$ scattering state verified¹⁴. In the KN -pentaquark system that we are concerned with, the ground state mass is quite high compared to that of one hadron. Furthermore, the excited states of the system, which are the discrete KN scattering states with N and K each with discrete lattice momentum in units of $2\pi/L$, are stacked up more compact than the radial excitation in the one hadron case when the lattice size is reasonably large (e. g. more than ~ 2.4 fm). Both of these factors require a large time separation or a more sophisticated fitting routine, such as fitting with Bayesian priors¹⁷ or with a variational approach, in order to obtain the spectral weights reliably⁴. The study of $I = 2$ $\pi\pi$ in Ref. [14] serves as a caveat for the authors' earlier work on pentaquark where they found the volume dependence of the spectral weight behaves like that of a one-particle state¹¹. Another calculation⁷, which observes little volume dependence of the spectral weight for the first excited state in the $1/2^-$ channel, employed variational method with two operators and used two exponentials to fit the spectral weights. As is widely known in fitting procedures, one cannot trust the results of the highest fitted state which is the second state in this case. As we pointed out in Sec. 2, the lesson learned from the variational calculations with multi-operators¹⁰ vs two operators² should be taken to heart. As such, the results of this study⁷ should be taken with a grain of salt.

Another way of distinguishing a scattering state from a pentaquark state is the clever idea of adopting the 'hybrid boundary condition'⁵ with the anti-periodic spatial boundary condition for the u and d quarks and periodic condition for the strange quark. This way, the energies of the KN scattering states will be raised compared to those from the usual periodic boundary condition; while those of the $uudd\bar{s}$ pentaquarks will stay the same⁵. Using this technique, it is found that the state near the KN threshold in the $1/2^-$ channel is a scattering state.

The third way to tell KN scattering state and pentaquark state apart is to examine the volume dependence of the spectrum^{10,7,11,14}. The one-particle state is not expected to be sensitive to the lattice volume when it is large enough for the specific quark mass; whereas, a weakly interacting two-particle state with relative momentum p will have a volume dependence since p is in units of $2\pi/L$. This approach applies well to our study since the KN interaction is weak and, as a result, it does not distort the discrete KN spectrum much from the non-interacting one. But it requires high statistics in order to discern the volume dependence which is milder compared to the $1/V$ scaling of the spectral weight.

4. Chiral Symmetry and Ghost States

It is learned from studying the quenched chiral logs in m_π, m_N ¹⁸, and the meson cloud effect in Roper resonance and $S_{11}(1535)$ ¹⁶ that chiral dynamics begins to play an important role in baryons when the pion mass is lower than ~ 300 MeV which characterizes the chiral regime. It is logically possible that the pentaquark state exists only for very light quark masses. After all, the first prediction of the anti-decuplet pentaquark is based on the chiral Skyrme model¹⁹. Since most of the lattice calculations are carried out outside this chiral regime, the conclusion drawn from these calculations may not be relevant in this case. To have a definite answer on the existence of the pentaquark will require calculations in the chiral regime with dynamical fermions. When in the chiral regime, one needs to be concerned with the presence of ghost states for the quenched approximation and the partially quenched case when the sea quark and valence quark masses do not match. The relevant ghost state in our case is $KN\eta'$ with the would be η' loop not fully developed to that of a physical η' . Instead, it has a double π pole which has the pion mass and breaks unitarity by giving a negative contribution to the 5-quark correlator. The ghost state is a practical problem in the $1/2^+$ channel in the chiral regime where the N, K and η' can be in relative S-wave which becomes lower than the P-wave KN state at certain mass and gives rise to a negative correlator⁴. In this case, the ghost state and the physical states have to be fitted together before drawing any conclusion¹⁶. After the $KN\eta'$ ghost state is removed, it is found that the lowest physical state is the KN P-wave scattering state in the $1/2^+$ channel⁴.

There is only one lattice calculation which claims to have detected a pentaquark in the $1/2^+$ channel⁹. When extrapolated to the chiral limit, it approaches the KN threshold. This contradicts the findings in all the other calculations. We speculate that this could be due to the contamination of the $KN\eta'$ ghost state. Even though the quark masses are somewhat higher than the chiral regime, the fact that the spatial volume is small in this study ($L = 1.8$ fm) makes it more susceptible to the negative contribution from the three-particle $KN\eta'$ ghost state which scales like $1/V^2$. Since the would be η' has the same mass as the pion, the ghost state is expected to reach the KN threshold at the chiral limit. Another noticeable feature of the calculation is that the overlap fermion results are found to be higher than those of the Wilson fermion for m_π heavier than ~ 550 MeV in the positive parity channel and ~ 700 MeV in the negative parity channel²⁰. In the $1/2^+$ case, the mass difference can be as large as ~ 300 MeV. This needs an explanation. In contrast, ground state masses in the $1/2^-$ channel from 6 different calculations with both the overlap and Wilson fermions are plotted in Fig. 28 in Ref. [8] and they pretty much lay on top of each other in the pion mass range from 420 MeV to 890 MeV.

5. Conclusions

To summarize, we have learned several lessons from studying the 5-quark system which includes two-particle scattering states in the spectrum. First of all, the

8 *Keh-Fei Liu*

appearance of the interpolation field does not necessarily betray the structure of the hadron. To learn about the structure of a hadron, it is better to study the three- and four-point correlation functions. Secondly, it is essential to discern the one- or two-particle nature of the observed state before making a claim of having detected a pentaquark. In order to resolve states close by, it is necessary to perform variational calculation with more interpolation fields than the number of states of interest. Lastly, it is necessary to go down to small enough quark masses where the chiral dynamics dominates and get rid of the quenched or partially quenched ghost state in the $1/2^+$ channel before one can be sure if the pentaquark exists.

After close examination of the existing lattice calculations, we conclude that there is no convincing evidence that any pentaquark state has been observed so far.

Of course, ‘absence of evidence is not evidence of absence’¹². To finally settle the issue, one needs to carry out realistic dynamical fermion calculations at physical quark masses, do a variational calculation to study the spectrum and the volume dependence of the spectral weight, and remove the potential ghost states in the partially quenched case.

Acknowledgments

This work is partially supported by USDOE grants DE-FG05-84ER40154 and DE-FG02-95ER40907. The authors thank C. Alexandrou, T. Draper, Z. Fodor, M. Oka, and S. Sasaki for useful discussions.

References

1. T. Nakano, plenary talk of this conference.
2. F. Csikor, Z. Fodor, S. D. Katz and T. G. Kovacs, *JHEP* **0311**, 070 (2003).
3. S. Sasaki, *Phys. Rev. Lett.* **93**, 152001 (2004).
4. N. Mathur, F.X. Lee, A. Alexandru, C. Bunnhold, Y. Chen, S.J. Dong, T. Draper, I. Horvath, K.F. Liu, S. Tamhankar, and J.B. Zhang *Phys. Rev.* **D70**, 074508 (2004).
5. N. Ishii, T. Doi, H. Iida, M. Oka, F. Okiharu and H. Suganuma, *Phys. Rev.* **D71**, 0340001 (2005).
6. C. Alexandrou, G. Koutsou and A. Tsapalis, *Nucl. Phys. B (Proc.Suppl.)* **140** 275 (2005).
7. T. T. Takahashi, T. Umeda, T. Onogi and T. Kunihiro, *Phys. Rev.* **D71**, 114509 (2005).
8. B. G. Lasscock, et al., *Phys. Rev.* **D72**, 014502 (2005).
9. T. W. Chiu and T. H. Hsieh, *Phys. Rev.* **D72** 034505 (2005)
10. F. Csikor, Z. Fodor, S.D. Katz, T.G. Kovacs, B.C. Toth, hep-lat/0503012.
11. C. Alexandrou and A. Tsapalis, hep-lat/0503013.
12. K. Holland and K.J. Juge, hep-lat/0504007.
13. O. Jahn, J. W. Negele, D. Sigaev, hep-lat/0509102..
14. C. Alexandrou and A. Tsapalis, hep-lat/0509139.
15. R. Jaffe and F. Wilczek, *Eur.Phys.J.* **C33**, S38 (2004).
16. N. Mathur, Y. Chen, S.J. Dong, T. Draper, I. Horvath, F.X. Lee, K.F. Liu, and J.B. Zhang, *Phys. Lett.* **B605**, 137 (2005), [hep-lat/0306199].
17. G.P. Lepage, et al. *Nucl. Phys. Proc.Suppl.* **B106**, 12 (2002); Y. Chen, S.J. Dong,

- T. Draper, I. Horvath, K.F. Liu, N. Mathur, S. Tamhankar, C. Srinivasan, F. X. Lee, J.B. Zhang, hep-lat/0405001.
18. Y. Chen, S.J. Dong, T. Draper, I. Horváth, F.X. Lee, K.F. Liu, N. Mathur, and J.B. Zhang, *Phys. Rev.* **D70**, 034502 (2004).
19. D. Diakonov, V. Petrov, and M.V. Polyakov, *Z. Phys.* **A359**, 305 (1997).
20. T. W. Chiu and T. H. Hsieh, hep-lat/0509175.